

On the long-term tidal evolution of GJ 436b in the presence of a resonant companion

Rosemary A. Mardling^{1*}

¹*School of Mathematical Sciences, Monash University, Victoria, 3800, Australia*

Accepted ... Received ...; in original form ...

ABSTRACT

In order to explain the significant orbital eccentricity of the short-period transiting Neptune-mass planet GJ 436b and at the same time satisfy various observational constraints and anomalies, Ribas, Font-Ribera and Beaulieu have proposed the existence of an eccentric low-mass companion planet at the position of the outer 2:1 resonance. The authors demonstrate the viability of their proposal using point-mass three-body integrations, arguing that as long as the system appears to be dynamically stable, the short-term secular variations ought to dominate the long-term dissipative evolution. Here we demonstrate that if one includes tidal dissipation, *both* orbits circularize after a few times the circularization timescale of the inner planet. We conclude that with or without a nearby companion planet, in or out of the 2:1 resonance, the Q -value of GJ 436b must be near the *upper* bound estimate for Neptune if the system is as young as 1 Gyr, and an order of magnitude higher if the system is as old as 10 Gyr. We show detail of passage through resonance and conclude that even out of resonance, a companion planet should still be detectable through transit timing variations.

Key words: planetary systems – celestial mechanics – methods: analytical – planetary systems: formation

1 INTRODUCTION

GJ 436b was discovered in a radial velocity survey by Butler et al (2004), and has since been observed in transit by Gillon et al (2007) and Alonso et al (2008) as the only transiting hot Neptune to date. Its system parameters are listed in Table 1 together with those for the hypothetical companion of Ribas, Font-Ribera & Beaulieu (2008). Here m and R are the mass and radius of the body (Deming et al 2007), a is the semimajor axis, e is the orbital eccentricity, ω , Ω and i are the argument of periastron, longitude of the ascending node and inclination respectively, the latter two measured relative to the line of sight, M is the mean anomaly, Q is the Q -value, k is the Love number and R_g is the radius of gyration. All angles are in degrees; round brackets refer to the hypothetical orbital data used by Ribas et al. (2008), while square brackets refer to initial data used in the simulation presented here (see Section 3.1). Note that the mass and radius of GJ 436b are respectively 1.35 and 1.094 times those of Neptune. As an extremely short-period system with a significant eccentricity, it has attracted considerable attention because it would appear that such a system should long ago have circularized. However, there is one factor in

particular which works against the intuition that GJ 436b ought to be circularized, and that is that the planet's distance from the star in units of its *radius* is quite large at 160 compared to, for example, HD 209458b at 76. Since the circularization timescale depends on the fifth power of this ratio, it turns out to be relatively long if one uses estimates for the Q -value of Neptune. The latter has been estimated by Banfield & Murray (1992) to be in the range $1.2 \times 10^4 < Q_N < 3.3 \times 10^5$, while Tittlemore & Wisdom (1989) estimate the Q -value of Uranus, whose mass is 0.85 times that of Neptune, to be less than 3.9×10^4 . Using the expression

$$\tau_{circ} = e_b / \dot{e}_b \simeq \frac{2}{42\pi} \left(\frac{Q_b}{k_b} \right) \left(\frac{m_b}{m_*} \right) \left(\frac{a_b}{R_b} \right)^5 P_b \quad (1)$$

for the circularization timescale of a synchronous system (Goldreich & Soter 1966),¹ where the subscript b refers to quantities associated with GJ 436b, m_* is the stellar mass and $P_b = 2.64$ d is the orbital period, one obtains the range $5.3 \times 10^7 \text{ yr} < \tau_{circ} < 1.5 \times 10^9 \text{ yr}$, where we have used the Banfield & Murray (1992) estimates for Neptune

* E-mail: mardling@sci.monash.edu.au

¹ Note that Goldreich & Soter (1966) use a modified Q -value, Q'_b , which absorbs the Love number such that $Q'_b = 3Q_b/2k_b$.


Table 1. Observed and hypothetical orbital and structural data for a GJ 436 two-planet system

	m	R	a (AU)	e	ω	Ω	i	M	Q	k	R_g/R
star	$0.452 M_\odot$	$0.452 R_\odot$							$[10^5]$	$[0.028]$	$[0.276]$
GJ 436b	$23.2 M_\oplus$	27,600 km	0.0287	0.15	343		86.54				
			[0.0295]	[0.2]	[343]	[90]	[86.54]	(0)	?	[0.346]	[0.511]
GJ 436c	$(4.7 M_\oplus)$	-	(0.045)	(0.2)	(265)	[30]	(96.54)	(107.6)	-	-	-

as well as Jupiter’s quadrupole Love number (0.34). Being estimated to lie in the range 1-10 Gyr, the age of the system is quite uncertain (Torres 2007). Thus in order for the non-zero eccentricity to be simply a result of a circularization time which is longer than the age of the system, this simple analysis suggests that Q_b must be greater than $2.3 \times 10^5 (\tau_{age}/\text{Gyr})$, where τ_{age} is the age of the system.

In this Letter we focus on the proposal of Ribas et al. (2008) that, given the circularization timescale is considerably less than the age of the system, the eccentricity is sustained by the presence of a low-mass companion planet positioned at the location of the outer 2:1 resonance, in an orbit which is inclined at around 10° to that of GJ 436b. The authors claim to have found a strong peak at 5.2 days in a periodogram analysis of the RV data, consistent with a body in the 2:1 resonance and with a false alarm probability of 20%. This is supported by a fit to the residuals of the two-body fit to the inner orbit. Moreover, the authors claim that harbouring a planet in the 2:1 resonance allowed them to produce the observed eccentricity of GJ 436b with a planet whose mass is low enough not to have previously been detected in the RV data.² Their scenario was further strengthened by the fact that it provides a natural explanation for the apparent change in the inclination to the line of sight of the orbit of GJ 436b, i_b , which would bring it into transit between the time of the null result of Butler et al (2004) and the positive detection of Gillon et al (2007). The authors estimated that a rate of change of i_b of around $0.1^\circ \text{ yr}^{-1}$ was needed to be consistent with these observations (note that this generally includes a combination of nutation *and* precession of the orbital plane). However, Alonso et al (2008) have more recently reported an upper limit for the current rate of change of i_b of $0.03 \pm 0.05^\circ \text{ yr}^{-1}$. Since $0 < di_b/dt < 0.03^\circ \text{ yr}^{-1}$ for about 20% of the precession cycle (see Figure 4(c)), this in itself does not rule out a Ribas et al-type system.

In the following section we briefly consider the tidal evolution of a single-planet system, putting a more accurate lower bound on Q_b for that case. In Section 3 we study the proposed orbital solution of Ribas et al. (2008) using a direct integration scheme for three bodies which includes perturbing accelerations due to tidal dissipation and spin-orbit coupling in both the star and the innermost planet, and the post-Newtonian relativistic contribution to the potential of the star (Mardling & Lin 2002). We demonstrate that long-term evolution tends towards the doubly-circular state, independent of the tidal circularization timescale of the outer planet, with the orbits inclined to each other by

**Figure 1.** Past and future evolution of GJ 436b without a companion. The system cannot be older than $0.38(Q_b/10^5)$ Gyr.

a fixed angle. Section 4 discusses stability while Section 5 presents a conclusion.

2 GJ 436B WITHOUT A COMPANION

The simplest explanation for the significant orbital eccentricity of GJ 436b is that the circularization timescale is longer than the age of the system. Figure 1 shows the results of integrating the secular equations (Mardling & Lin 2002) for a single synchronized planet backwards and forwards in time, with system parameters taken from Table 1 and $Q_b = 10^5$ (solid curves). The equations used are based on the analyses of Hut (1981) and Eggleton, Kiseleva & Hut (1998) in which a constant time lag is assumed for the tidal bulge, and are correct for any eccentricity (in as far as the constant time lag assumption is correct). If one assumes that no forces other than those included in the simulation were acting in the past, and also that the Q -values of the planet and star as well as their radii have not changed, these results suggest that the system cannot be older than $0.38(Q_b/10^5)$ Gyr. This is in contrast to the results obtained if one integrates equations which are correct to second-order only in the eccentricity (dashed curves). The latter approximation is clearly adequate for estimating the current circularization timescale, but is entirely inadequate for understanding the past evolution. In particular, the second-order analysis does not allow an upper bound estimate of the age of the system. While alternative tidal models such as those which assume constant lag angles for all tidal components

² Note that we find a maximum value of only 0.1 for e_b over a modulation cycle using their initial conditions, independent of initial phases and longitudes.

(Goldreich & Soter 1966), or those which include the dynamical tide for high eccentricity (Mardling 1995) may produce slightly different functional dependence on the eccentricity were the analysis to be carried out, they are not likely to change the estimated age of the system by much.

Other forces are likely to have been operating early in the life of the system, including those due to the rapid rotation of the star, larger stellar and planetary radii, the presence of a protoplanetary disk, coupling of the stellar and planetary magnetic fields (Laine, Lin & Dong 2008), and different Q -values. Such factors will have determined the “initial” orbital state of the system, that is, the state of the system when such influences became negligible, and it seems likely that the eccentricity would not have been much larger than its present value if the single-planet scenario is correct.

The main conclusion one can draw from this simple analysis is that if GJ 436b does not have companions capable of sustaining the observed eccentricity, the upper bound on its age in terms of Q_b puts a *lower* bound on the Q -value of GJ 436b of $Q_b > 3 \times 10^5 (\tau_{age}/\text{Gyr})$, a value slightly higher than that gleaned from equation 1. In particular, it is approximately equal to the *upper* bound for the Q -value of Neptune if the system is as young as 1 Gyr, and an order of magnitude larger if it is as old as 10 Gyr.

3 LONG-TERM TIDAL EVOLUTION OF TWO-PLANET SYSTEMS

The second simplest explanation for a non-zero eccentricity in a system like GJ 436 is the presence of a companion planet or star. In general the induced eccentricity of a stable system will vary quasi-periodically, with periods of variation dominated by the rate of change of the angle between the apsidal lines, the rate of change of the resonance angle(s) if the system is in a resonance, and the rate of change of the *argument* of periastron if the system is significantly non-coplanar (Murray & Dermott 2000). The amplitude of variation depends on the ratios of semimajor axes and planet masses as well as the initial eccentricities and the angle between the apsidal lines, and when relativistic effects are important, on the ratio of the inner planet mass to the stellar mass. Formulae for the amplitude and period of variation of the eccentricities are given in Mardling (2007) for coplanar non-resonant systems with moderate inner eccentricity ($e_b \lesssim 0.2$). The expression for the amplitude is also accurate for moderately non-coplanar and/or resonant systems, while that for the modulation period severely overestimates the true value for a system in or near the 2:1 resonance because for such a close system, higher order and resonant terms should be included.

A mistake often made is that secular variations of the orbital elements persist for the lifetime of the system, even when dissipative forces are significant (eg. Maness et al. 2007). In fact such variations are damped out after a few circularization timescales, with the system evolving to a pseudo-equilibrium configuration which itself evolves on a generally longer timescale. A secular theory for two-planet coplanar systems with dissipation has recently been developed (Mardling 2007), in which it is shown that the eccentricity of the innermost planet together with the angle between the apsidal lines, η , evolve towards a fixed point in




Figure 2. Possible values for m_c and e_c corresponding to $e_b^{(eq)} = 0.15$ with $a_c = 0.045$ AU. No solutions exist for $e_c < 0.18$, while essentially *all* of those with $e_c > 0.18$ are unstable.

(e_b, η) space on a timescale of three times the circularization timescale of the inner planet, after which the system evolves to the doubly circular state on timescale given by equation (60) in Mardling (2007). *Note that this timescale is independent of the Q -value of the outermost planet.* When resonant terms are important and/or the system is moderately inclined, the behaviour is only slightly modified (as long as the system is stable). The fixed point or equilibrium eccentricity, a quantity which is independent of the initial values of e_b , phases and longitudes, is given by

$$e_b^{(eq)} = \frac{(5/4)(a_b/a_c) e_c \varepsilon_c^{-2}}{\left| 1 - \sqrt{a_b/a_c} (m_b/m_c) \varepsilon_c^{-1} + \gamma \varepsilon_c^3 \right|}, \quad (2)$$

where $\varepsilon_c = \sqrt{1 - e_c^2}$ and $\gamma = 4(n_b a_b/c)^2 (m_*/m_c)(a_c/a_b)^3$, with n_b the mean motion of the inner planet and c the speed of light.³ Note, however, that (2) was derived assuming that the average value of the outer eccentricity doesn’t vary much on the circularization timescale. When it does (as happens for the hypothetical GJ 436 system because the outer mass is so low), the estimate (2) tends to *overestimate* the equilibrium eccentricity. Thus (2) can be regarded as an upper bound for $e_b^{(eq)}$.

3.1 GJ 436b with a resonant companion

If the circularization timescale of GJ 436b is less than a third of the age of the system, it will already have reached and evolved past the equilibrium eccentricity. Thus we begin by calculating the range of companion masses and eccentricities capable of producing an equilibrium eccentricity of 0.15, that is, the observed eccentricity of GJ 436b. As in the previous Section, this will allow us to put a lower bound on its Q -value. Using this value in (2) as well as a period ratio of 2, we can rearrange the equation to write m_c as a function of e_c . The result is plotted in Figure 2 which shows that no solutions exist for $e_c < 0.18$. Solutions corresponding to $e_c \gtrsim 0.18$ are actually unstable (see next section) so that in fact *no* systems exist for which the equilibrium eccentricity is equal to the observed eccentricity. (In fact, the equilibrium

³ The quantity γ is the ratio of the relativistic to the three-body contribution to the rate of apsidal advance of the innermost planet.

eccentricity corresponding to the hypothetical companion of Ribas et al. (2008) (with $e_c = 0.2$) is 0.06.)

One can conclude from this that if GJ 436b *does* have a single nearby low-mass companion in or near the 2:1 resonance, the system cannot yet have evolved to its pseudo-equilibrium state. Its circularization timescale must therefore be longer than one third of the age of the system, thereby providing the weaker constraint on the Q -value of GJ 436b than in the single-planet case that $Q_b > 7.7 \times 10^4 (t_{age}/\text{Gyr})$. In fact, we can do better than this, and argue that after only one circularization timescale the system will be in apsidal libration with zero minimum e_b and a maximum which depends on e_c (Mardling 2007). From stability considerations e_c can't be much more than around 0.3 (given it needs to be able to move safely through the resonance as it tidally evolves; see Figure 5). Since this corresponds to a *maximum* value of e_b in a libration cycle of 0.12, we must have that $\tau_{circ} > \tau_{age}$, putting the same lower bound on Q_b as in the single-planet case.

We finish this section by demonstrating the behaviour discussed above. Using the data in Table 1, we performed a direct integration using the code described in the Introduction (Mardling & Lin 2002). The star's Q -value is an estimate, while the Love numbers k (twice the apsidal motion constant) and radii of gyration R_g for the star and planet correspond to $n = 3$ and $n = 1$ polytropes respectively (Sterne 1941), the latter often taken to approximate the structure of Jupiter. The spin period of the star was taken to be 20 days while the spin of the planet was taken to be synchronous with the orbital motion. Both were taken to be aligned with the orbit normal.

The secular analysis in Mardling (2007) demonstrates that varying the Q -value of the inner-most planet merely changes the timescale on which the system evolves towards the final state while not affecting the local secular oscillation period. Since tidal dissipation in the planet dominates the tidal evolution while the inner orbit's eccentricity is non-zero, we took $Q_b = 0.1$ in order to show the detail of various stages of evolution, and confirmed that the evolutionary timescale scales linearly with Q_b for $Q_b = 1$ and 10. We also confirmed that the system remains stable for 2×10^5 yr with $Q_b = 10^6$, consistent with the results of Section 3.2. Figure 3 shows 4000 years of evolution, equivalent to $4 (Q_b/10^5)$ Gyr for general Q_b , except that $Q_b/0.1$ as many secular oscillations will have occurred in that time. It passes through its present (hypothetical) configuration after $1600(Q_b/0.1)$ yr. Panel (a) shows the evolution of e_b (bottom curve) and e_c , with the system entering resonance at around $200(Q_b/0.1)$ yr, and entering apsidal libration (panel (b)) when e_b (temporarily) hits zero at $1000(Q_b/0.1)$ yr. The latter occurs on a timescale of $1 \tau_{circ}$ after which the system evolves towards the quasi-equilibrium phase on a timescale of $2 \tau_{circ}$. The subsequent approach of both eccentricities to zero is well approximated by the secular theory and occurs on a timescale of around $6 \tau_{circ}$ in this case (equation (60) Mardling 2007). Panel (c) shows the approach to a fixed value of around 10.3° of the relative inclination.

Figure 4 shows detail of the passage through resonance. While the amplitude of variation of the eccentricity is only slightly enhanced (so that estimates provided by the secular theory are reasonably accurate), the amplitude of variation of the orbital period ratio P_c/P_b suddenly increases as the




Figure 3. Approach to the doubly circular state of a resonant two-planet GJ 436 system. See text for discussion.




Figure 4. Detail of the resonant phase of Figure 3. Note that the period and amplitude of these variations is independent of Q_b . See text for discussion.

system crosses the separatrix and enters the 2:1 resonance, with a width of around 0.035. The corresponding variation of the orbital period of GJ 436b is around 15 mins, a significant increase on the variation outside the resonance, whose effect on the transit timing would be easily measured (note that the libration period for this system is around 200 days). For




Figure 5. A stability map of the region surrounding the 2:1 resonance for a range of values of e_c .

coplanar systems in the 2:1 resonance with $m_b, m_c \ll m_*$, this variation is given by (Mardling, in preparation)

$$\begin{aligned} \delta P_b/P_b &= 2 \left[1 + (m_b/m_c) \sigma^{2/3} \right]^{-1} \delta \sigma / \sigma \\ &\simeq 2^{-2/3} (m_c/m_b) \delta \sigma, \end{aligned} \quad (3)$$

where $\sigma = P_o/P_i$, $\delta \sigma = \sigma - 2$, that is, the “distance” from resonance, and the approximation holds for $m_c \ll m_b$.⁴ However, the GJ 436 system passes through the resonance in around $300(Q_b/0.1) \text{ yr} = 0.3(Q_b/10^5) \text{ Gyr}$ so unless Q_b is at least equal to the upper estimate of Neptune’s Q -value, we would not expect to see the system in resonance now (given that it was deposited into the resonance around the time of formation). Alonso et al (2008) find no obvious departure from linear ephemeris and conclude that the proposed resonant solution of Ribas et al. (2008) is unlikely. Once the system leaves the resonance, the amplitude of variation of P_b is given by (3) with $\delta \sigma$ replaced by the quantity $\delta \sigma - \sqrt{\delta \sigma^2 - \Delta \sigma^2}$, where $\Delta \sigma$ is the width of the resonance (Mardling, in preparation). δP_b reduces quickly as the system crosses the resonance, however, even when the system becomes doubly circularized δP_b is still significant at around one minute. Thus we conclude that even out of resonance, the companion planet proposed by Ribas et al. (2008) would be detectable through transit timing variations given current accuracies (Alonso et al 2008).

Panel (c) shows the variation of the inclination of GJ 436b to the line of sight, i_b . The resonance has very little effect on the period and amplitude of variation, with the average rate of change of i_b equal to approximately $0.06^\circ \text{ yr}^{-1}$. This compares to the observational upper bound of $0.03 \pm 0.05^\circ \text{ yr}^{-1}$ by Alonso et al (2008).

3.2 Stability

Figure 5 shows a stability map for a system composed of GJ 436b and a companion planet with a mass of $4.7 M_\oplus$, aligned periastra, a mutual inclination of 10° , and with initial companion eccentricity e_c and period $P_c = \sigma P_b$ indicated by the position in the plot. Three-body integrations were performed, with stability being determined using the

procedure described in Mardling (2008). Initial conditions corresponding to unstable systems are indicated in the figure by red dots; stable systems are left blank. A clearly defined boundary is evident, indicating that a system with a sufficiently low initial value of e_c would be free to tidally evolve through the 2:1 resonance without the danger of instability. Other resonances are also evident, for example, the 3:2 and the 5:2.

4 CONCLUSION

The main conclusion from this study is that with or without a single nearby companion planet, in or out of the 2:1 resonance, the Q -value of GJ 436b must be greater than the upper bound estimate for Neptune if the age of the system is around 1 Gyr, and up to an order of magnitude greater for an age of up to 10 Gyr. Passage through resonance of a Ribas et al. two-planet system occurs on a timescale of $0.3(Q_b/10^5) \text{ Gyr}$, remaining stable throughout and beyond. However, even if it were now no longer in resonance, the companion planet should still be detectable through transit timing variations.

REFERENCES

- Agol E., Steffen J., Sari R., Clarkson W., 2005, MNRAS, 359, 567
- Alonso R., Barbieri M., Rabus M., Deeg H. J., Belmonte J. A., Almenara, J. M., 2008, arXiv:0804.3030
- Banfield D., Murray N., 1992, Icarus, 99, 390
- Butler R. P., Vogt S. S., Marcy G. W., Fischer D. A., Wright J. T., Henry G. W., Laughlin G., Lissauer J. J., 2004, ApJ, 617, 580
- Deming D., Harrington J., Laughlin G., Seager S., Navarro S. B., Bowman W. C., and Horning K., 2007, ApJL, 667, L199
- Eggleton P. P., Kiseleva L. G., Hut, P. 1998, ApJ, 499, 853
- Gillon M., et al., 2007, A&A, 472, L13
- Goldreich P., Soter S., 1966, Icarus, 5, 375
- Holman M. J., Murray N. W., 2005, Science, 307, 1288
- Hut P., 1981, A&A, 99, 126
- Laine R. O., Lin D. N. C., Dong S., 2008, arXiv:0804.0975
- Maness H. L., Marcy G. W., Ford E. B., Hauschildt P. H., Shreve A. T., Basri G. B., Butler R. P., Vogt S. S., 2007, PASP, 119, 90
- Mardling R. A., 1995, ApJ, 450, 722
- Mardling R. A., 2007, MNRAS, 382, 1768
- Mardling R. A., 2008, in Aarseth, S. J., Tout, C. A., Mardling, R. A., eds, The Cambridge N-Body Lectures, Lecture Notes in Physics, Springer, in press
- Mardling R. A., Lin D. N. C., 2002, ApJ, 573, 829
- Murray C. D., Dermott S. F., 2000, Solar System Dynamics. Cambridge Univ. Press, Cambridge, UK
- Ribas I., Font-Ribera A., Beaulieu J.-P., 2008, ApJL, 677, L59
- Sterne T. E., 1941, Proceedings of the National Academy of Science, 27, 93
- Tittlemore W. C., Wisdom, J., 1989, Icarus, 78, 63
- Torres G., 2007, ApJL, 671, L65

⁴ See also Holman & Murray (2005) and Agol et al (2005) but note that our expression is independent of the eccentricities, and is an approximation to an expression valid for any masses.